

International Journal of Applied Environmental Sciences

Daily Flight Activity of *Trigona laeviceps* AND *T.*
minangkabau in Red Pepper (*Capsicum annum* L.)
Plantations in Low and High Lands of West Sumatra

Dewirman Prima Putra, Dahelmi, Siti Salmah and Etti Swasti

DAILY FLIGHT ACTIVITY OF *Trigona laeviceps* AND *T. minangkabau* IN RED PEPPER (*Capsicum annuum* L.) PLANTATIONS IN LOW AND HIGH LANDS OF WEST SUMATRA

Dewirman Prima Putra¹, Dahelmi², Siti Salmah², Etti Swasti³

¹Departement of Agronomy, Faculty of Agriculture, Ekasakti University, Padang 25163, Indonesia

²Departement of Biology, Faculty of Mathematics and Natural Science, Andalas University, Padang 25163, Indonesia

³Departement of Agronomy, Faculty of Agriculture, Andalas University, Padang 25163, Indonesia.

corresponding author: de_wirman_pp@yahoo.com

Abstract

Red pepper is vegetable commodity which is primary plant in West Sumatra. It can be found every where from low to high lands. Stingless bees are species of insect that visit red pepper plantations. Objective of research was to study daily flight activity of two species of stingless bees, *T. laeviceps* and *T. minangkabau* in red pepper plantations in low and high lands. Stingless bees were captured from red pepper plantations using scan sampling method from 07.00 a.m until 04.00 p.m. Each observation was done for 15 minutes per hour per day per location. *Trigona laeviceps* and *T. minangkabau* were dominant species found, each with 492 individuals (58.16%) and 354 individuals (41.84%). *T. laeviceps* was found in ten locations of red pepper plantations, eight were in high lands. *T. minangkabau* was found in seven locations, four were in high lands. Daily visit of *T. laeviceps* and *T. minangkabau* in red pepper plantations was from 08.00 a.m until 03.00 a.m. The peak of visits of both occurred at 11.00 p.m. *T. laeviceps* was captured in altitudes 50 – 910 m asl. at temperature 22-35.5 °C and relative humidity (RH) 32-78 %. Locations of *T. minangkabau* captured was at altitudes 30–900 m asl. at temperature 20-35 °C, and RH 37-78 %. Population of *T. laeviceps* in high lands was higher than in low lands. In other hand, population of *T. minangkabau* in high lands was the same as in low lands. The presence of *T. laeviceps* and *T. minangkabau* in red pepper plantations was from 08.00 a.m until 3.00 p.m. The peak daily activity of both for food foraging occurred at 11.00 a.m.

Key words : Stingless bees, *Capsicum annuum*, low land, high land, west Sumatra

1. INTRODUCTION:

Red pepper (*Capsicum annuum* L.) can be cultured either in low or high lands, in rice field land or dry land [1]. In West Sumatra red pepper is a vegetable crop having important advantages, and even in Minangkabau society it is considered as primary crop. Therefore red pepper plantations are distributed in many districts and town in West Sumatra which are the centres of red pepper productions.

Red pepper flowers are chasmogamy, pollination occurs at the time of flowers are opened. Thus, in red pepper there is possibility for cross pollination to occur [2]. This is the reason why the production of red pepper per plant is higher when planted in group than individually. The increase of yield due to cross pollination was 7.6-36.8% [3].

Stingless bees are insects visiting red pepper plantations in West Sumatra. The species were dominated by *Trigona laeviceps* (55.85 %) and *T. minangkabau* (40.18 %) [4]. Both species could be found in low and high lands red pepper plantations.

The presence of *T. laeviceps* and *T. minangkabau* as pollinating insects in red pepper plantations could increase fruit set 9.66 % and 12.32 %, number of seeds 45.91% and 56.36 %, and number of fruits per plant 25.06 % and 29.31% [5]. Objective of research was to study daily flight activity of *T. laeviceps* and *T. minangkabau* in red pepper plantations in low and high land of West Sumatra.

2. MATERIALS AND METHODS

The captures of stingless bees were done in red pepper production centers located in high lands (> 500 m asl) and low lands (< 200 m asl) in West Sumatra. High land locations were: Jampang Manganti (1) and Koto Panjang (2) Lima Puluh Kota District, By Pass (3) and Baringin (4) Payakumbuh Town, Ganting (5) and Gunung (6) Padang Panjang Town, Sabu (7) and Sikaladi (8) Tanah Datar District, Kubang Putihah (9) and Batu Taba (10) Agam District, Panyangkalan (11) and Batu Banyak (12) Solok District. Low land locations were:

Data Analysis

Data on species and number of individuals of stingless bees captured in each location of red pepper plantation per time period were presented descriptively. Stingless bees captured and climatic factors (temperature and relative humidity) were display in tables and graphs.

3. RESULTS AND DISCUSSION

Two species of stingless bees were found visiting red pepper plantations in production centres in West Sumatra. They were *Trigona laeviceps* with 492 individuals (58.16%) and *T. minangkabau* with 354 individuals (41.84 %). *T. laeviceps* and *T. minangkabau* were captured more in highlands than in the lowlands.

Trigona laeviceps was found in ten locations of red pepper plantations. Eight were in high lands (Koto Panjang Suliki and Japang Manganti in Lima Puluh Kota District, By Pass and Baringin in Payakumbuh, Ganting and Gunung in Padang Panjang, Sabu in Tanah Datar, and Batu Banyak in Solok) with 399 individuals (81,10 %). Two locations were in low lands (Kampung Dalam in Padang and Kubang in Bayang Pesisir Selatan) with 93 individuals (18.90%).

T. minangkabau was found in seven locations, four in high lands (Sikaladi in Tanah Datar, Kubang Putihah and Batu Taba in Agam, and Panyangkalan in Solok) with 198 individuals (55.93%). Three locations were in low lands (Lubuk Minturun in Padang, Ketapiang and Lubuk Alung in Padang Pariaman) with 156 individuals (44.07%). The number of individuals of *T. laeviceps* and *T. minangkabau* captured in various locations was presented in Table 1.

In high lands 66.67% of *T. laeviceps* and 33.33% of *T. minangkabau* were captured. While in low lands they were 40% of *T. laeviceps* and 60 % of *T. minangkabau*. *T. laeviceps* was captured in altitudes 50 – 910 m asl. at temperature 22-35.5 °C and relative humidity (RH) 32-78 %. Locations of *T. minangkabau* captured was at altitudes 30–900 m asl. at temperature 20-35 °C, and RH 37-78 %.

Daily visit of *T. laeviceps* and *T. minangkabau* to red pepper plantations was from 08.00 a.m until 03.00 p.m. The peak of visit of both occurred at 11.00 a.m. (106 individuals of *T. laeviceps* at $30.2 \pm 2,4$ °C and RH $46.9 \pm 5,9$ %, and 85 individuals of *T. minangkabau* at $30 \pm 2,6$ °C and RH $49.57 \pm 5,1$ %). There was no visit made by the two species before 08.00 a.m and after 3.00 p.m.

Food foraging of *T. laeviceps* occurred at temperature of locations 22.8-32.75 °C and RH 71.1-40.9 % and those of *T. minangkabau* 23.29-32.78 °C and RH 73.14-44.01%. Population density of *T. laeviceps* and *T. minangkabau* in daily food foraging in high and low land at temperature and RH of red pepper plantations was presented in Figure 2 and 3.

Trigona laeviceps is one species of stingless bees living in tropical and subtropical regions [9]. Daily activity of stingless bees is influenced by climate like temperature, RH, and light intensity [10,11,12]. Three hundred ninty nine individuals (81.10 %) of *T. laeviceps* were captured in high lands and 93 individuals (18,90%) in low lands. The higher percentage of *T. laeviceps* capture in high land was assumed to be caused by most of red pepper production centres in West Sumatra were located in high land (>500 m asl). Out of 12 locations of *T. laeviceps* capture, eight locations (66.67%) were high lands and out of five locations of *T. laeviceps* capture, two of them (40 %) were low land. Salmah *et al.*[13] reported that 53.8% of *T. laeviceps* were found in low lands and 37.5% in high lands. However, the center of red pepper plantations are often found in the highlands. thus *T. laeviceps* also mostly found in the highlands. Temperature at research locations in high lands ranged 22-35.5 °C and in low lands was 21-35 °C. Even though stingless bees can live at 34-36°C [14], however *T. laeviceps* preferred cool areas of high land.

Different from *T. laeviceps*, *T. minangkabau* captured was 198 individuals (55.93 %) in high lands and 156 individuals (44.07 %) in low lands. It indicated that the activity of *T. minangkabau* was not influenced by altitude. This condition could be seen at percentage of

capturing locations of *T. minangkabau*, out of five capturing locations three of them were low lands (60.00 %) and out of 12 locations, four of them were high lands (33.33 %). Salmah *et al.* [13] reported that 55.8 % of *T. minangkabau* (include *T. minangkabau* f. darek) were found in low lands and 43.8 % in high lands.

The presence of *T. laeviceps* and *T. minangkabau* in red pepper plantations was from 08.00 a.m. until 3.00 p.m. The peak daily activity of both for food foraging occurred at 11.00 a.m (*T. laeviceps* 106 individuals and *T. minangkabau* 85 individuals) (Fig. 2 and 3). This case relates to the occurrence of red pepper flowers to blossom which happen in the morning (07.00-09.00 a.m) and corollas usually open in the first 3 hours after the sun rises [15]. The same result was also reported by Putra *et al.* [16] that the peak of daily activity of *Apis cerana* for food foraging in red pepper plantations occurred at 11.00 a.m and of *T. laeviceps* occurred from 11.00-12.00 a.m. However, the activity of *A. cerana* started from 07.00 a.m and *T. laeviceps* started from 08.00 a.m. Al Abbadi [17] reported that the peak daily activity of honey bees in eggplants happened from 09.00-10.00 a.m at temperature 28-33⁰C, and of Bombus bee from 08.00-09.00 a.m at 25-28⁰C local time. The same condition also happened on red pepper and bell pepper plants, the peak daily activity of honey bees occurred from 09.00-10.00 a.m and of Bombus bee from 08.00-09.00 a.m. Jordan time.

The higher number of *T. laeviceps* captured in high lands and *T. minangkabau* in low lands was caused by size and color of these both *Trigona* were little different. *T. minangkabau* had body length 3.2 – 3.5 mm, metasoma was predominantly chesnut brown, while *T. laeviceps* had body length 4-4.5 mm and blackish brown body color. Willmer and Corbet [18] stated that small and light color of stingless bees were more resistant to full sun shine than those having big size and dark body color.

The insects visit to flowers decreased by the decrease of nectar volume and the increase of temperature. At 12.00-13.00 a.m the temperature was at its culminated peak, while RH was at

its lowest condition and therefore the weather was very dry. This conditions caused the volume of nectar decreased even though its concentration increased due to evaporation of nectar solution. Besides that, the reduce of stingless bees activity for food foraging in day time was caused by the reduce of food stock. Roubik *et al.* [19] reported that the concentration of nectar sugar ranged from 20-67%. The optimum concentration of nectar sugar was 40-50%. Further, Biesmeijer *et al.* [20] stated that concentration of nectar sugar increased from 07.00 until 11.00 a.m and it was stable at 11.00 a.m.

Gojmerac [21] stated that when temperature increased the energy needed for flying to search food was higher, while the amount of nectar available in nature as energy source was little. Therefore, in their activity for food foraging, stingless bees considered the potential energy that could be obtained. If energy spent for food foraging was higher than energy obtained the stingless bees would decrease their activity. In the morning the availability of nectar was higher so that the energy obtained was higher too compared to in a day time. Therefore, in a day time stingless bees activity decreased.

4. CONCLUSION

T. laeviceps was captured in locations at altitudes 50 – 910 m asl and that of *T. minangkabau* was at altitudes 30–900 m asl. Population of *T. laeviceps* at high lands was higher than in low lands, in other hand the population of *T. minangkabau* at high lands was the same as in low lands. The presence of *T. laeviceps* and *T. minangkabau* in red pepper plantations was from 08.00 a.m. until 3.00 p.m. The peak daily activity of both for food foraging occurred at 11.00 a.m.

ACKNOWLEDGEMENT

It is my pleasure to acknowledge Directorate General of Higher Education Ministry of Research and Technology-Higher Education and Kopertis Area X for providing research fund

on Hibah Disertasi Doktor. My thanks also goes to Language Centre of Andalas University for the proofreading of this article.

REFERENCES

- [1] Sumarni, N., and Muharam A., 2005, Cultivation of Chili Pepper, Vegetable Crops Research, Lembang-Bandung.
- [2] Pickersgill, B., 1997, "Genetic resources and breeding of *Capsicum* spp," Euphytica, 96(1), 129-133.
- [3] Greenleaf, W.H., 1986, "Pepper Breeding in Breeding of Vegetable Crops," M. Bassett (ed.). AVI Publishing Company, INC, pp. 67-134.
- [4] Putra D. P., Dahelmi, Salmah S., and Swasti E., 2016a, "Species Diversity of Stingless Bees (Hymenoptera: Meliponini) in Chili Pepper (*Capsicum annum* L.) Plantation in West Sumatra," Int. J. Sci. Res., 5(4), 1527-1532
- [5] Putra D. P., Dahelmi, Salmah S., and Swasti E., 2016b, "Pollination in chili pepper (*Capsicum annum* L.) by *Trigona laeviceps* and *T.minangkabau* (Hymenoptera, Meliponini)," J. Entomol. Zool. Studies, 4(4), 191-194.
- [6] Martin, P., and Bateson P., 1993, "Measuring behaviour an introductory guide," Cambridge University Press, Cambridge, UK, Chap. 6.
- [7] Sakagami S. F., Inoue T., and Salmah S., 1990, "Stingless bees of Central Sumatra in Natural history of social wasps and bees in equatorial Sumatra," S.F., Sakagami *et al.*, eds., Hokkaido University Press, Sapporo, Japan, pp. 125–137.
- [8] Michener, C., 2007, "The bees of the world," Johns Hopkins University Press, Baltimore, Maryland, USA. pp. 803-829.
- [9] Sakagami, S. F., Yamane, S., and Hambali, G. G., 1983, "Nest of some southeast Asian Stingless bee," Bull. Fac. Educ. Ibaraki Univ. Nat. Sci., 32,1-21.
- [10] Contrera, F.A.L., Imperatriz-Fonseca, V.L., and Nieh, J.C., 2004, "Temporal and climatological influences on flight activity in the stingless bee *Trigona hyalinata* (Apidae, Meliponini)," Rev. Tecn. Amb., 10(2), 35-43.
- [11] Souza, B.A., Carvalho, C.A.L., and Alves, R.M.O., 2006, "Flight activity of *Melipona asilvai* Moure (Hymenoptera: Apidae)," Braz. J. Biol., 66(2B), 731-737.
- [12] Nascimento, A.S., Pereira, L.L, Carvalho, C.A.L., Machado, C.S., Souza, M.O., and Souza, B.A., 2012, "Flight activity of the eusocial bee *Melipona quadrfasciata anthidioides* (Hymenoptera: Apidae, Meliponini)," Magistra, 24(5), 112-118.
- [13] Salmah, S., Inoue, T., and Sakagami, S. F., 1990, "An analysis of apid bee richness (Apidae) in Central Sumatra in Natural history of social wasps and bees in equatorial Sumatra," S.F., Sakagami *et al.*, eds., Hokkaido University Press, Sapporo, Japan, pp. 139-174.
- [14] Amano, K., 2004, "Attempts to introduce stingless bees for the pollination of crops in greenhouse conditions in Japan," Laboratory of Apiculture, National Institute of Livestock and Grassland Science Tsukuba, Ibaraki 305-0901, Japan.
- [15] Aleemullah, M., Haigh A. M., and Holford, P., 2000, "Anthesis, anther dehiscence, pistil receptivity and fruit development in the Longum group of *Capsicum annum*," Aust. J. Exp. Agric., 40(5), 755–762.

- [16] Putra, R. E., Permana, A.D., and Kinasih, I., 2014, "Application of Asiatic Honey Bees (*Apis cerana*) and Stingless Bees (*Trigona laeviceps*) as Pollinator Agents of Hot Pepper (*Capsicum annuum* L.) at Local Indonesia Farm System," Hindawi Publishing Corporation. Psyche. Volume 2014, Article ID 687979, 5 pages.
- [17] Al-Abbadi, S. Y. A., 2009, "Efficiency of different pollination treatments on solanaceae yields grown in plastic house," J. Biol. Sci., 9(5):464–469.
- [18] Wilmer, P.G., and Corbet, S.A., 1981, "Temporal and microclimatic partitioning of the floral resources of *Justicia aurea* amongst a concourse of pollen vectors and nectar robbers," Oecologia, 51(1), 67-78.
- [19] Roubik, D.W., Yanega, D., Aluja, S. M., Buchmann, S. L, and Inouye, D. W., 1995, "On optimal nectar foraging by some tropical bees (Hymenoptera: Apidae)," Apidologie, 26(3), 197-211.
- [20] Biesmeijer, J.C., Smeets, M., Richter, J.A.P., and Sommeijer, M.J., 1999, "Nectar foraging by stingless bees in Cost a Rica: botanical and climatological influences on sugar concentration of nectar collected by *Melipona*," Apidologie, 30(1), 43–55.
- [21] Gojmerac, W, L., 1983, "Bee, Beekeeping honey And Pollonation," Avi Publishing Company, Inc., Westport, Connecticut.

Table 1. Number of individuals of *T. laeviceps* and *T. minangkabau* captured on various red pepper locations in high and low lands in West Sumatra

No	Location	Altitude (m)	<i>T. laeviceps</i> (individual)	<i>T. minangkabau</i> (individual)
High lands				
1.	Japang Manganti	520	50	-
2.	Koto panjang Suliki	650	49	-
3.	By Pass Payakumbuh	600	62	-
4.	Baringin	600	37	-
5.	Gantiang	740	51	-
6.	Gunung	760	54	-
7.	Sabu Tanah Datar	910	53	-
8.	Sikaladi Tanah Datar	880	-	60
9.	Kubang Putih Agam	900	-	40
10.	Batu Taba Agam	900	-	49
11.	Batu banyak Solok	800	43	-
12.	Panyangkalan Solok	830	-	49
Total			399 (81,10 %)	198 (55,93 %)
Low lands				
1.	Kampung Dalam Kota Padang	50	57	-
2.	Lubuk Minturun Kota Padang	30	-	61
3.	Kubang Bayang Pessel	100	36	-
4.	Ketaping Padang Pariaman	30	-	44
5.	Lubuk Alung Padang Pariaman	40	-	51
Total			93 (18,90 %)	156 (44,07 %)
Total number of individuals			492	354




Figure 2. Population density of *T. laeviceps* in daily food foraging at temperature and RH of red pepper plantations at low land (LL) and high land (HL).




Figure 3. Population density of *T. minangkabau* in daily food foraging at temperature and RH of red pepper plantations at low land (LL) and high land (HL).




Dewirman Prima Putra <de_wirman_pp@yahoo.com>
Kepada: submit@ripublication.com

Sel, 6 Jun 2017 jam 23.07 ☆

Dear submission desh International Journal of Applied Environmental Science
I have sent to our manuscript entitled, "DAILY FLIGHT ACTIVITY OF *Trigona laeviceps* and *T. minangkabau* IN RED PEPPER (*Capsicum annuum* L.) PLANTATIONS IN LOW AND HIGH LANDS OF WEST SUMATRA on May 22, 2017.
Therefore I request to confirm whether my paper has been received and how next.
Best regards

Dewirman Prima Putra



Journal DPP.docx
242kB

RIP Research India Publications

Head Office: B-2/84, Ground Floor, Rohini Sector-16, Delhi-110089 INDIA

Phone no.: +91-11-65394240 Fax No.: +91-11-27297815

Website: www.rippublication.com Email: ripublication@vsnl.net

COPYRIGHT TRANSFER FORM

Journal Title: _____

Paper Code: _____

I/(We) confirm that the enclosed article entitled:

Authored by: 1. _____ 2. _____
3. _____ 4. _____

(This form is signed by the corresponding author on behalf of all of them)

has not previously been published in whole or in part, is not currently being considered elsewhere for publication, and, if accepted for publication in the above Journal, will not be published elsewhere in any language, without the consent of the editor and the publisher.

2) I acknowledge that it is a condition of acceptance by the editor that the publisher, Research India Publications acquires automatically the copyright in the manuscript throughout the world.

3) I confirm that I have obtained all the necessary permissions to include in the paper items such as quotations, figures, and the results of government sponsored research.

4) I enclose where necessary written permission of authors and publishers to use any copyright material (e.g. previously published figures and tables).

Author 1

Signature: _____ Name: _____ Date: _____

Institution: _____ Dept: _____

Street: _____ City: _____ State: _____

Zip: _____ Country: _____ Country Code: _____

Phone: _____ Fax: _____ Email: _____

Author 2

Signature: _____ Name: _____ Date: _____

Institution: _____ Dept: _____

Street: _____ City: _____ State: _____

Zip: _____ Country: _____ Country Code: _____

Phone: _____ Fax: _____ Email: _____

Author 3

Signature: _____ Name: _____ Date: _____

Institution: _____ Dept: _____

Street: _____ City: _____ State: _____

Zip: _____ Country: _____ Country Code: _____

Phone: _____ Fax: _____ Email: _____

Author 4

Signature: _____ Name: _____ Date: _____

Institution: _____ Dept: _____

Street: _____ City: _____ State: _____


Zip: _____ Country: _____ Country Code: _____

Phone: _____ Fax: _____ Email: _____


Phone: _____ Fax: _____ Email: dahelmi@gmail.com

(This form is signed by the corresponding author on behalf of all of them)

Author 3

Signature:  Name: Siti Salmah Date: 22 Juni 2017
Institution: Faculty of Mathematics and Natural Science, Andalas University Dept: Biology
Street: Limau Manih City: Padang State: Sumatera Barat
Zip: 25163 Country: Indonesia Country Code:
Phone: _____ Fax: _____ Email: salmah.idrus@gmail.com

Author 4

Signature:  Name: Etti Swasti Date: 22 Juni 2017
Institution: Faculty of Agriculture, Andalas University Dept: Agronomy
Street: Limau Manih City: Padang State: Sumatera Barat
Zip: 25163 Country: Indonesia Country Code:
Phone: _____ Fax: _____ Email: ettiswasti14@yahoo.com



Research India Publications <ripjournal@gmail.com>
Kepada: Dewirman Prima Putra



Rab, 21 Jun 2017 jam 17.40 ☆

Dear **Dr. Dewirman Prima Putra,**

Paper Code: 56829

We are very pleased to inform you that your paper "**DAILY FLIGHT ACTIVITY OF *Trigona laeviceps* AND *T. minangkabau* IN RED PEPPER (*Capsicum annuum* L.) PLANTATIONS IN LOW AND HIGH LANDS OF WEST SUMATRA**" is accepted by our Editor-in-chief for our journal **International Journal of Applied Environmental Sciences (IJAES)**. Please find attached copyright transfer form.

Publication charges is \$250.00 (US \$ Two Hundred and Fifty only).

Publication Charges includes One Print Journal copy, PDF Copy and free online access to journal issue. Please send us signed copy of copyright form along with **CHEQUE/DRAFT in US\$ PAYABLE TO "RESEARCH INDIA PUBLICATIONS"**. You can also send payment by Credit Card.

id-Racti@itu-subar@mail

Dear **Dr. Dewirman Prima Putra**

We are very pleased to inform you that we are now registered with Crossref so now we can get the DOI number of our published journal articles. Your paper is published in our Journal "**International Journal of Applied Environmental Sciences (IJAES) Vol.12 No.08 (2017)**". If you want to get the DOI number of your paper with a DOI publication certificate you have to pay **\$25** for your paper. After payment you will receive the final PDF file of your paper with DOI number with publication certificate.

You can send us fee of \$25 by Credit Card or bank transfer.

If pay by bank transfer, please add an extra US\$ 5.00 for bank charges. Our bank information is given below

SWIFT CODE -----CITIINBX
Credit to -----RESEARCH INDIA PUBLICATIONS
Account No. -----0031384222
Name of Bank-----CITIBANK N. A
Address -----27, Central Market, West Punjabi Bagh, NEW DELHI-110026, INDIA

Please inform us when you transfer payment to our account so that we can track your payment

Thanking you

With kind regards,

Viveka Nand

=====



Dewirman Prima Putra <de_wirman_pp@yahoo.com>
Kepada: ripublication@vsnl.net

Kam, 6 Jul 2017 jam 12.54 ☆

I already payment of publishing a journal to the bank CITIBANK N.A with the title paper "DAILY FLIGHT ACTIVITY OF *Trigona laeviceps* AND *T. minangkabau* IN RED PEPPER (*Capsicum annum* L.) PLANTATIONS IN LOW AND HIGH LANDS OF WEST SUMATRA" (PAPER CODE #: RIP-56829- IJAES).
Along with this I also created COPYRIGHT TRANSFER FORM from author

Thank & Regards

Dewirman Prima Putra


[Unduh semua lampiran sebagai file zip](#)



Payment.jpg
253.1kB



Payment 2.jpg
92.8kB



Form.jpg
196.7kB



Form 001.jpg
87.1kB